

# Asymmetry of Jet $k_T$ in Longitudinal Polarized p+p Collisions in PHENIX at RHIC

R. Hobbs
For the PHENIX collaboration
University of New Mexico




## Outline

- Measuring transverse momentum of partons in the proton
- Looking for a correlation of this measurement with spin direction
- Initial measurements in Run03
- Status of Run05
- Summary




# **PH** ENIX $k_{\rm T}$ , $j_{\rm T}$ from azimuthal correl.




# PHIENIX


## Origin of $k_{\rm T}$

$$\frac{1}{2} \langle p_T^2 \rangle_{pair} ? \langle k_T^2 \rangle \qquad ?$$






An example - J/? production.



 $p_{T}$ ?<sub>J/?</sub> =1.8?0.23?0.16 GeV/c Phys. Rev. Lett. 92, 051802, (2004).




# Hard Scattering $k_{\rm T}$



transverse

- acoplanar in P<sub>I</sub> ? P<sub>T</sub> space
- collinear in P<sub>X</sub> ? P<sub>Y</sub> space




transverse


- acoplanar in P<sub>I</sub> ? P<sub>T</sub> space
- acoplanar in P<sub>X</sub> ? P<sub>Y</sub> space




# k<sub>T</sub> from Orbital Motion

One can consider the possibility that spin-correlated transverse momentum (orbital angular momentum) may contribute to jet  $k_T$ .






e.g., Meng Ta-chung et al., *Phys. Rev. D 40 (1989)* 



unlike-sign helicities






#### Azimuthal correlation function


$$C_{ij}(?\,\emph{?})\,?\,\textit{norm}\,? \frac{dN^{real}_{\phantom{real}ij}}{d?\,\emph{?}_{ij}}\,/\,\frac{dN^{mixed}_{\phantom{mixed}ij}}{d?\,\emph{?}_{ij}}$$

?° - h? correlation functions


#### Not corrected for acceptance



#### Corrected for acceptance



Jet kinematics



$$p^2_{\text{out}}? = p^2_{\text{Ta}} \sin^2 ? ? ? ?$$

$$p_{\text{out}}^2 = p_{\text{Ta}}^2 \sin^2 ? ? ? 2? k_{\text{Ty}}^2 ? 2? k_{\text{Ty}}^2 ? 2? k_{\text{Ty}}^2 ? 2? k_{\text{Ty}}^2 ? 2 k_{\text{h}}^2 x_{\text{h}}^2 x$$

$$x_{
m h} = p_{
m Ta}/p_{
m Tt}$$

$$\sqrt{\langle j_T^2 \rangle} ? \sqrt{2} \frac{p_{Tt} p_{Ta}}{\sqrt{p_{Tt}^2 ? p_{Ta}^2}} ?_N$$

$$\begin{array}{c|c}
p_{\text{Tt}} & j_{\text{T}} \\
\hline
p_{\text{Ta}} & j_{\text{T}} & \hat{q}
\end{array}$$

$$\hat{x}_h^{?1}\langle z_t\rangle\sqrt{\langle k_T^2\rangle}$$
 ?  $x_h^{?1}\sqrt{\langle p_{out}^2\rangle?\langle j_{Ty}^2\rangle(1?x_h^2)}$ 


partonic 
$$\hat{x}_h$$
 ?  $\frac{\left\langle \hat{p}_{Ta} \right\rangle}{\left\langle \hat{p}_{Tt} \right\rangle}$ 


hadronic 
$$x_h$$
 ?  $\frac{p_{Ta}}{p_{Tt}}$ 



## Spin Sorted Analysis

 Do exactly the same analysis sorted on same and opposite helicity bunch crossings, extract <zkt><sub>RMS</sub> and look at the difference.





Oct. 27, 2005


9




### Run03 Data

Like sign


Unlike sign






trigger  $p^0$ 1<  $p_{Tt}$ <3 Gev/c 3< $p_{Tt}$ <7 Gev/c

Associated  $h^?$  1<  $p_{Ta}$ <2.5 GeV/c








# Systematic Check

- Helicity assignments are randomized, and then the k<sub>T</sub> difference calculated for each randomized set.
- The width of the distribution for all the randomized sets should be the same as our statistical errors on the previous plot.






## Run03 Data

Its too early to make a definite statement about the apparent excess as the systematic uncertainties are being evaluated.

However, there is an ongoing analysis of 10x more stat. and 2x better polarization in run $05 \approx$  should yield a definite answer.





- We have an analysis tool that allows us to measure  $k_{\mathsf{T}}$  initial state transverse momentum of partons.
- We are studying this effect in longitudinal spin-sorted collisions to see if there is a spin-dependent coherent component of  $k_{\rm T}$ .
- Is there a connection to parton OAM?
   Theoretical guidance needed!

#### Outlook

- Run05 has ~x10 statistics, so that the uncertainty reduced by factor 2-3.
- It has ~x2 in polarization, so the effect grows by ~x4.